Hyperglycemia at hospital admission is associated with severity of the prognosis in patients hospitalized for COVID-19.

Alberto Coppelli¹, Rosa Giannarelli¹, Michele Aragona, Giuseppe Penno^{1,2}, Marco Falcone², Giusy Tiseo², Lorenzo Ghiadoni², Greta Barbieri², Fabio Monzani², Agostino Virdis², Francesco Menichetti², and Stefano Del Prato^{1,2} on behalf of the Pisa-COVID-19 Study Group*

¹Section of Diabetes and Metabolic Diseases, Azienda Ospedaliero-Universitaria Pisana, Pisa, Italy,

²Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy,

*The Pisa COVID-19 Study –

PISA Covid-19 Study Group: Agostini o Degl'Innocenti Sabrina¹, Antognoli Rachele², Baldassarri Rubia³, Bertini Pietro³, Biancalana Martina¹, Borselli Matteo¹, Brizzi Giulia³, Calsolario Valeria², Carrozzi Laura⁴, Celi Alessandro⁴, Cinotti Francesco⁵, Cipriano Alessandro⁵, Corradi Francesco³, De Marco Salvatore⁶, Della Rocca Alessandra³, Forfori Francesco³, Forotti Giovanna⁶, Galfo Valentina⁷, Guarracino Fabio³, Malacarne Paolo³, Masi Stefano⁶, Mengozzi Alessandro⁶, Monfroni Marco³, Morea Alessandra⁷, Nencini Elia¹, Park Naria⁵, Paterni Simone², Piagnani Chiara³, Ruberti Francesca⁶, Santini Massimo⁵, Sciuto Maria⁶, Spinelli Stefano¹.

¹Emergency Medicine Unit, Azienda Ospedaliera Universitaria Pisana; ²Geriatrics Unit, Azienda Ospedaliera Universitaria Pisana; ³Department of Anaesthesia and Intensive Care, Azienda Ospedaliera Universitaria Pisana; ⁴Respiratory Unit, Azienda Ospedaliera Universitaria Pisana; ⁵ Emergency Department, Azienda Ospedaliera Universitaria Pisana; ⁶Internal Medicine Department, Azienda Ospedaliera Universitaria Pisana, ⁷Infectious Diseases Unit, Azienda Ospedaliera Universitaria Pisana; Pisa (Italy).

Supplementary Table 1. Clinical and laboratory parameters in the three groups at hospital admission. The number in parenthesis provides the number of observations.

	All	Glycemia < 7.78 mmol/L	Glycemia >7.84 mmol/L	Diabetes	p
No., %	271	149, 55%	66, 24.3%	56, 21%	
Age, years	72 [58-82]	63 [53-79]	75 [63-82] *	75.5 [69-84] *	0.02
Male/Female	181/90	93/56	49/17	39/17	NS
BMI, Kg/m ²	26.2 [23.4-29.3]	25.7 [23.4-28.9]	27.3 [24.6-31.3]	26.9 [23.2-28.5]	NS
Glycemia, mmol/L	7.11 [5.71-8.57]	5.99 [5.38-6.72]	8.57 [8.18-10.47] *	9.18 [7.67-12.71] *	0.002
HbA1c, mmol/mol - %	44 [38-52] – 6.2 [5.6-6.9]	36 [36-38.5] – 5.4 [5.4-5.7]	43.5 [40-47] – 6.1 [5.8- 6.5]*	49 [42.5-60.5] – 6.6 [6.0-7.7] *	0.03
Total cholesterol, mmol/L	4.29 [3.02-4.86]	4.19 [2.91-4.97]	3.80 [2.81-5.23]	4.45 [3.22-4.78]	NS
HDL cholesterol, mmol/L	0.96 [0.68-1.25]	0.96 [0.81-1.14]	0.68 [0.55-1.07]	1.09 [0.86-1.46]	NS
LDL cholesterol, mmol/L	2.34 [55-126]	90 [1.43-3.28]	2.42 [52-3.25]	2.24 [1.79-3.28]	NS
Triglycerides, mmol/L	3.20 [2.21-4.00]	3.30 [2.63-3.90]	3.28 [2.00-4.91]	3.09[2.08-4.24]	NS
Statin therapy	46/267 [17.2]	19/149 [12%]	7/66 [10%]	20/56 [36.5%] *	<0.05
Systolic blood pressure, mmHg	130 [120-144]	130 [120-140]	130 [117-145]	126 [114-144]	NS
Diastolic blood pressure, mmHg	77 [67-85]	80 [70-85]	73 [65-82]	75 [65-82]	NS
Anti-hypertensive therapy	132/166 [50%]	47/149 [31.5%]	22/66 [33.3%]	43/56 [76.8%] *	<0.05
Creatinine, μmol/L	88.4 [70.7-114.9]	84.9 [69.8-104.3]	97.2 [70.7-123.8]	106.1 [79.6-159.1] *	0.01
eGFR, ml/min	75.1 [48.1-91.5]	79.5 [59.1-95.2]	73.1 [38.9-87.7]	65.1 [34.6-81.7] *	0.01
White blood cell, x10 ⁹ per L	6.7 [5.1-9.6]	6.4 [5.0-9.1]	7.5 [5.1-10.6]	6.7 [5.6-11.1]	NS
White blood cell, >11 x10 ⁹		23/146 [16%]	14/65 [21%] *	14/55 [25%] *	0.005
Neutrophil, x10 ⁹ per L	5.0 [3.4-8.0]	4.6 [3.2-7.2]	5.8 [3.7-8.7] *	4.9 [4.0-8.8]	<0.05
Lymphocyte, x10 ⁹ per L	0.95 [0.7-1.3]	1.0 [0.7-1.4]	0.7 [0.5-1.2] *	0.8 [0.7-1.3]	<0.05
Prothrombin, %	74 [64-82]	74 [64-82]	72 [64-77]	72 [63-85]	NS
Prothrombin <80%		84/124 [68%]	42/54 [78%] *	32/48 [67%]	<0.05
D-Dimer, mg/L	0.48 [0.26-0.87]	0.41 [0.22-0.67]	0.63 [0.37-1.28] *	0.63 [0.28-2.04] *	0.04
Fibrinogen, mg/dl	586 [425-730]	544 [405-726]	647 [506-730]	635 [409-793]	NS
Fibrinogen >400		87/114 [76%]	41/47 [87%] *	32/41 [78%]	<0.05

mg/dl					
Ferritin, mg/L [13-150]	642 [363-1213]	590 [376-1103]	780 [550-1345]	579 [339-1127]	NS
Ferritin >150 mg/L]		61/72 [85%]	25/26 [96%]	25/29 [86%]	NS
PCT, ng/ml	0.14 [0.07-0.38]	0.11 [0.06-0.21]	0.21 [0.11-0.69] *	0.19 [0.09-0.44] *	0.02
CRP, mg/dl	6.69 [2.69-14.04]	4.99 [1.92-11.86]	10.6 [4.29-16.67] *	11.4 [4.46-18.26] *	0.001
IL-1, pg/ml	1.4 [1.0-2.0]	1.4 [1.0-2.2]	1.4 [1.0-2.0]	1.4 [0.9-2.1]	NS
IL-6, pg/ml	20.1 [9.9-52.3]	14.9 [8.3-43.3]	27.4 [11.6-91.7] *	23.6 [12.2-65.1]	0.03
IL-10, pg/ml	3.6 [1.1-6.9]	2.6 [0.9-6.1]	5.5 [1.5-9.7] *	4.4 [2.6-8.7] *	0.03
TNF, pg/ml	8.2 [3.8-12.4]	7.2 [3.5-12.0]	9.2 [4.3-13.5]	9.3 [3.4-12.7]	NS
MCP, pg/ml	597 [398-833]	604 [385-793]	578 [386-831]	660 [467-1046]	NS
pO ₂	70 [58-88]	74 [61-90]	67 [56-85]	61 [55-85]	NS
PaO ₂ /FiO ₂	275 [195-338]	295 [221-367]	227 [140-290] *	262 [201-290]	<0.05

Data are presented as median [IQR], or n/N [%]. Data were compared with the Kruskal-Wallis test for continuous variables and chi-square test for categorical variables.

Supplementary Table 2. Univariate and multivariate Cox regression models for mortality in the overall population (271 patients) stratified in 3 categories (NG, normoglycemia; HG, hyperglycemia; DM, diabetes mellitus).

	Univariate HR (95% CI)	p	Multivariate Model 1 HR (95% CI)	P	Multivariate Model 2 HR (95% CI)	p	Multivariate Model 3 HR (95% CI)	P
Age, years	1.06 (1.04-1.08)	0.0001	1.07 (1.04-1.09)	0.0001	1.07 (1.04-1.10)	0.0001	1.07 (1.03-1.10)	0.0001
Male sex	1.48 (0.86-2.57)	0.159	2.07 (1.16-3.68)	0.013	2.25 (1.10-4.60)	0.070	1.70 (0.76-3.78)	0.194
Baseline glucose categories		0.017		0.088		0.064		0.038
<i>Normoglycemia</i>	1		1		1		1	
<i>Diabetes</i>	1.73 (0.92-3.25)	0.086	1.07 (0.56-2.04)	0.828	0.83 (0.36-1.89)	0.656	0.78 (0.29-2.09)	0.623
<i>Hyperglycemia</i>	2.19 (1.27-3.81)	0.005	1.80 (1.05-3.15)	0.040	1.96 (1.00-3.85)	0.049	2.39 (1.10-5.19)	0.028
Obesity (BMI >30 kg/m²)	2.80 (0.97-8.10)	0.055						
Smokers vs non-smokers	0.74 (0.37-1.47)	0.380						
Hypertension	2.22 (1.33-3.70)	0.002			1.21 (0.63-2.30)	0.564	1.02 (0.49-2.15)	0.954
Dyslipidemia	1.50 (0.84-2.67)	0.170						
Coronary artery disease	1.70 (0.95-3.03)	0.720						
Congestive heart failure	1.90 (0.94-3.82)	0.070						
Cerebrovascular disease	1.95 (1.02-3.74)	0.044			1.50 (0.50-4.53)	0.468	1.78 (0.55-5.75)	0.336
Cognitive impairment	2.38 (1.15-4.93)	0.020			1.27 (0.40-4.02)	0.680	1.25 (0.34-4.68)	0.738
COPD	1.95 (1.04-3.67)	0.038			0.76 (0.33-1.73)	0.511	0.84 (0.34-2.06)	0.700
CKD	2.56 (1.37-4.80)	0.003			0.68 (0.21-2.17)	0.512	1.34 (0.37-4.89)	0.655
Chronic liver disease	1.28 (0.46-3.56)	0.640						
Cancer	1.09 (0.59-2.00)	0.790						
Leucocytes, x10⁹/L	1.00 (1.00-1.00)	0.530						
Neutrophils, x10⁹/L	1.00 (1.00-1.00)	0.760						
Lymphocytes, x10⁹/L	0.99 (0.99-1.00)	0.008					1.00 (0.99-1.00)	0.208
PCT, ng/ml	1.01 (0.97-1.06)	0.600						
RCP, mg/dL	1.02 (0.99-1.04)	0.220						
D-Dimer, mg/L	1.04 (0.97-1.08)	0.810						
Prothrombin, %	0.98 (0.97-100)	0.032					0.99 (0.98-1.02)	0.716

<i>IL-6</i> , pg/mL	1.00 (1.00-1.01)	0.621
<i>IL-10</i> , pg/mL	1.00 (1.00-1.01)	0.109
<i>TNF</i> , pg/mL	1.00 (1.00-1.01)	0.070
<i>MCP</i> , pg/mL	2.56 (1.37-4.80)	0.283

Supplementary Table 3. Univariate and multivariate Cox regression models for mortality in the whole population (271 patients) stratified by quintiles of glucose at baseline.

	Univariate HR (95% CI)	p	Multivariate Model 1 HR (95% CI)	p	Multivariate Model 2 HR (95% CI)	p	Multivariate Model 3 HR (95% CI)	p
Age, years	1.06 (1.04-1.08)	0.0001	1.07 (1.05-1.10)	0.002	1.08 (1.05-1.11)	0.0001	1.08 (1.00-1.12)	0.0001
Male sex (vs female)	1.49 (0.86-2.58)	0.159	2.26 (1.26-4.04)	0.006	2.39 (1.15-4.95)	0.019	1.70 (0.74-3.89)	0.209
Quintiles of glucose		0.019		0.020		0.005		0.014
Q1	1		1		1		1	
Q2	1.37 (0.49-3.85)	0.550	1.39 (0.49-3.92)	0.533	1.79 (0.49-6.51)	0.377	1.50 (0.37-6.08)	0.569
Q3	1.83 (0.69-4.88)	0.227	2.17 (0.81-5.68)	0.121	4.34 (1.32-14.3)	0.016	3.87 (1.04-14.43)	0.044
Q4	3.57 (1.46-8.76)	0.005	3.33 (1.34-8.27)	0.009	5.98 (1.83-17.4)	0.003	5.91 (1.73-20.19)	0.005
Q5	2.32 (0.91-5.96)	0.079	1.40 (0.54-3.63)	0.488	1.71 (0.53-5.56)	0.370	1.70 (0.49-5.90)	0.402
Obesity (BMI >30 kg/m²)	2.80 (0.97-8.10)	0.055						
Smokers vs non-smokers	0.74 (0.37-1.47)	0.380						
Hypertension	2.22 (1.33-3.70)	0.002			1.16 (0.62-2.18)	0.642	1.00 (0.48-2.08)	0.995
Dyslipidemia	1.50 (0.84-2.67)	0.170						
Coronary artery disease	1.70 (0.95-3.03)	0.720						
Congestive heart failure	1.90 (0.94-3.82)	0.070						
Cerebrovascular disease	1.95 (1.02-3.74)	0.044			1.77 (0.67-4.65)	0.247	2.05 (0.72-5.85)	0.180
Cognitive impairment	2.38 (1.15-4.93)	0.020			1.16 (0.39-3.44)	0.787	1.15 (0.33-3.99)	0.828
COPD	1.95 (1.04-3.67)	0.038			0.94 (0.42-2.11)	0.887	0.98 (0.40-2.42)	0.973
CKD	2.56 (1.37-4.80)	0.003			0.72 (0.23-2.19)	0.559	1.09 (0.32-3.65)	0.893
Chronic liver disease	1.28 (0.46-3.56)	0.640						
Cancer	1.09 (0.59-2.00)	0.790						
Leucocytes, x10⁹/L	1.00 (1.00-1.00)	0.530						
Neutrophils, x10⁹/L	1.00 (1.00-1.00)	0.760						

Lymphocytes , x10 ⁹ /L	0,99 (0.99-1.00)	0.008	1.00 (0.99-1.00)	0.219
PCT , ng/ml	1.01 (0.97-1.06)	0.600		
RCP , mg/dL	1.02 (0.99-1.04)	0.220		
D-Dimer , mg/L	1.04 (0.97-1.08)	0.810		
Prothrombin , %	0,98 (0.97-1.00)	0.032	0.99 (0.97-1.02)	0.517
IL-6 , pg/mL	1.00 (1.00-1.01)	0.621		
IL-10 , pg/mL	1.00 (1.00-1.01)	0.109		
TNF , pg/mL	1.00 (1.00-1.01)	0.070		
MCP , pg/mL	2.56 (1.37-4.80)	0.283		

Supplementary Table 4. Treatments received and clinical outcomes

	Total n. 271	Normoglycemia <7.78 mmol/L n. 149	Hyperglycemia ≥7.78 mmol/L n. 66	Known diabetes n. 56	p
Treatments					
ICU Care, n [%]	80 [29.5]	36 [24.2]	30 [45.4] *	14 [25.0]	0.002
Invasive Mechanical Ventilation, n [%]	46 [17.0]	17 [11.4]	22 [33.3] *	7 [12.5]	0.002
Outcomes					
ARDS, n [%]	124 [45.8]	49 [32.9]	39 [59.1] *	36 [64.3] *	<0.05
Secondary infection, n [%]	22/198 [11.1]	15/114 [13.2]	3/45 [6.7]	4/39 [10.3]	NS
Hospitalization Duration [days]	14 [7-24]	13 [7-23]	16 [8-26]	13 [8-23]	NS
Mortality, n [%]	67 [24.7]	25 [16.8]	26 [39.4] *	16 [28.6]	0.002

Data are presented as number [percentages], or median [IQR], or n/N [%].

* p at least <0.05 vs Normoglycemia

Supplementary Figure 1. ROC analysis showing glucose level cut-off for mortality