Online Supplemental File

Roman Vangoitsenhoven, Rickesha Wilson, Deepa V Cherla, Chao Tu, Sangeeta R Kashyap, David E Cummings, Philip R Schauer, Ali Aminian. **Presence of Liver Steatosis is Associated with Greater Diabetes Remission after Gastric Bypass Surgery.**

Table S1. Patient characteristics according to presence of steatosis

	Steatosis (n=407)	No steatosis (n=112)	P-value
Age at time of surgery (years)	51 [44;57]	54 [49;61]	0.019
BMI before surgery (kg/m ²)	44,7 [40;50]	47,0 [42.1;53.1]	0.002
Duration of diabetes (years)	5 [2;10]	7 [3.7;13]	0.004
Preoperative insulin use			
No	264 (64,9%)	67 (59,8%)	0.35
Yes	143 (35,1%)	45 (40,2%)	
Diabetes medications	2 [1;3]	2 [1;3]	0.36
Preoperative HbA1c (%)	7,4 [6.5;8.6]	6,7 [6.1;8.0]	0.019
Poor glycemic control (HbA1c ≥7%)			0.003
No	155 (38,1%)	60 (53,6%)	<0.001
Yes	252 (61,9%)	52 (46,4%)	
Lobular inflammation			<0.001
No	100 (25,6%)	60 (53,6%)	
Yes	307 (75,4%)	52 (46,4%)	<0.001
Hepatocyte ballooning			
No	230 (56,5%)	105 (93,8%)	<0.001
Yes	177 (43,5%)	7 (6,3%)	
Fibrosis			<0.001
No	215 (52,8%)	93 (83,0%)	
Yes	192 (47,2%)	19 (17,0%)	0.019
Remission of diabetes			
No	196 (48,2%)	68 (60,7%)	0.019
Yes	211 (51,8%)	44 (39,3%)	

Values are n (%) or median [interquartile range]