The effects of B1344, a novel fibroblast growth factor 21 analog, on nonalcoholic steatohepatitis in nonhuman primates

Aoyuan Cui^{1,5}, Jian Li^{2,5}, Shaohui Ji^{3,5}, Fengguang Ma¹, Genbei Wang^{2,4}, Yaqian Xue¹, Zhengshuai Liu¹, Jing Gao¹, Jun Han², Ping Tai⁴, Tony Wang³, Jianxun Chen³, Xiaohui Ma^{4*}, Yu Li^{1*}

¹CAS Key Laboratory of Nutrition, Metabolism and Food Safety, Shanghai Institute of Nutrition and Health, Shanghai Institutes for Biological Sciences, University of Chinese Academy of Sciences, Chinese Academy of Sciences, Shanghai 200031, China; ²Tasly Biopharmaceuticals Co., Ltd, Shanghai 201203, China; ³Kunming Biomed International, Yunnan 650500, China; ⁴Research Center of Pharmacology and Toxicology, Tasly Academy, Tasly Pharmaceutical Co. Ltd, Tianjin 300410, China;

⁵These authors contribute equally to this work;

Supplemental Fig. 2

Cui A, Li J, Ji S, et al.

Supplemental Fig. 3

Cui A, Li J, Ji S, et al.

Supplemental Figure Legends

Fig.S1. Effects of B1344 on body weight in MCD diet fed mice. Eight-week-old male C57BL/6 mice were fed on MCD diet for 2 weeks, and then treated with B1344 (0.125 mg/kg or 0.2 mg/kg) or vehicle by subcutaneous injection for 2 weeks (short term injection) or 4 weeks (long term) as indicated. (A) Body weight was determined weekly. (B) mRNA levels of FGFR1 and β Klotho in livers of mice treated with B1344 were determined by real-time PCR. The data are represented as the mean $\pm$ SEM, n=7-9.

Fig.S2. Energy intake and plasma metabolic parameters in cynomolgus monkeys treated with B1344. 40 obese monkeys that met the enrolment criteria were treated with 0.5 mg/kg B1344 twice weekly (BIW), 1 mg/kg B1344 BIW, 2 mg/kg B1344 BIW, 2 mg/kg B1344 once weekly (QW) or vehicle by subcutaneous injection for 11 weeks, and then monitored for additional 4 weeks washout period. (A-B) Plasma LDL cholesterol (A) and HDL cholesterol (B) levels are represented as changes from baseline. n=6-8.

Fig.S3. B1344 improves glucose metabolism in cynomolgus monkeys. (A-C) Metabolic parameters were measured at the indicated days and represented as changes from baseline, including fasted blood glucose (FBG) (A), hemoglobin A1c (HbA1c) (B) and fasted insulin (C). (D) intravenous glucose tolerance test (IVGTT) was performed on day 45 since the first injection. (E) Glucose AUC measurements calculated from IVGTT. (F) Plasma insulin was measured during the IVGTT. (G) Insulin AUC measurements were calculated. n=6-8. *p<0.05, versus the vehicle.

Table S1. Quantitative RT-PCR primers

Gene	Species	Forward primer	Reverse primer
CD68	mouse	CATCCCCACCTGTCTCTCTC	TTGCATTTCCACAGCAGAAG

F4/80	mouse	ACCACAATACCTACATGCACC	AAGCAGGCGAGGAAAAGATAG
IL-1 β	mouse	TTCGTGAATGAGCAGACAGC	GGTTTCTTGTGACCCTGAGC
MCP1	mouse	CAGCCAGATGCAGTTAACGC	GCCTACTCATTGGGATCATCTTG
TNF α	mouse	CGTCAGCCGATTTGCTATCT	CGGACTCCGCAAAGTCTAAG
Col3a1	mouse	CTGTAACATGGAACTGGGGAAA	CCATAGCTGAACTGAAAACCACC
TGF β	mouse	CCTGCAAGACCATCGACATG	GCGAGCCTTAGTTTGGACAG
TIMP	mouse	CTCAAAGACCTATAGTGCTGGC	CAAAGTGACGGCTCTGGTAG
ACTA2	mouse	ATGCTCCCAGGGCTGTTTTCCCAT	GTGGTGCCAGATCTTTTCCATGTCG
FGFR1	mouse	CACCAAGAAGAGCGACTTCC	CATGGATGCACTGGAGTCAG
β Klotho	mouse	GATGAAGAATTCCTAAACCAGGT	AACCAAACACGCGGATTTC
β -actin	human	GATGAGATTGGCATGGCTTT	GTCACCTTCACCGTTCCAGT

Supplemental Experimental Procedures

Expression and purification of B1344. B1344, the FGF21 analog, is produced in E. coli Rosetta (DE3) as described previously¹. The bacteria clones are subjected to experiments such as shake flask expression and screening, fermenter expression, and plasmid retention rate, to screen and obtain the clones of the strain for establishing the seed lot. A three-level seed lot comprising the primary seed lot, master seed lot and working seed lot is established, and the seed lot is subjected to systemic verification and the results complies with the laws and guidelines of National Medical Products Administration of China (NMPA).

The fermentation is performed via a three-stage culturing process, including primary shake flask culture, secondary amplification in a 50 L-sized seed fermenter, followed by fermentation and manufacturing in a 500 L-sized fermenter. No animal-derived components are added throughout the

entire fermentation and manufacturing process. The yield of the target protein is mainly present in the form of soluble proteins, avoiding additional treatment such as renaturation of inclusion bodies. Next, a complete set of drug substance purification processes is performed to evaluate the purity, yield and bioactivity of the target protein, followed by removal of impurities such as residual host DNA and HCP, so as to finally determine the purification process based on affinity chromatography and using Capto Q and Capto Q ImpRes as the primary steps of fine purification. The target protein is subjected to site-directed N-terminal PEG modification, fine purification via Capto Q, ultrafiltration/diafiltration, and sterile filtration to finally obtain recombinant protein B1344.

Reagents and antibodies—B1344 was expressed and purified as described previously¹. Lyophilized human recombinant FGF21 (cat. 41189) that is dissolved in deionized water to prepare the working stock solution was obtained from Antibody and Immunoassay Services at the University of Hong Kong, China. Antibodies against pERK1/2 (cat. 9106) and ERK1/2 (cat. 9102) were purchased from Cell Signaling Technology (Beverly, MA). Horseradish peroxidase-conjugated anti-mouse (cat. sc-2005) and anti-rabbit (cat. sc-2004) secondary antibodies were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Alanine/Aspartate Transaminase Determination Kit from Shensuo UNF (Shanghai, China) were used for measurement of levels of ALT and AST in plasma of mice.

Intravenous glucose tolerance test (IVGTT) in the monkey—IVGTT was performed on day 45 since the first injection. Monkeys were fasted overnight and anesthetized continuously with ketamine hydrochloride (5-10 mg/kg), 0.5 g/kg glucose was dosed intravenously in 30 seconds. Vein blood samples were collected at the indicated time points and examined by automatic biochemistry analyzer.

Histological analysis of the NASH CRN Score. NASH CRN Score were evaluated according to the NASH Clinical Research Network (CRN) scoring system^{2, 3}. The NAFLD activity score (NAS) ranging from 0 to 8 were defined as the sum of the scores for steatosis (0-3): 0<5%, 1: 5-33%, 2: 34-66%, 3: >66%; lobular inflammation (0-3): 0: none, 1: <2 foci/20x field, 2: 2-4 foci/20x field, 3: >4 foci/20x field; hepatocellular ballooning (0-2): 0: none, 1: mild/few; 2: moderate-marked/many. Fibrosis (0-4) were evaluated: 0: none, 1: perisinusoidal or periportal; 1A: mild, zone 3, perisinusoidal; 1B: moderate, zone 3, perisinusoidal; 1C: portal/periportal; 2: perisinusoidal and portal/periportal; 3: bridging fibrosis; 4: cirrhosis.

Pharmacokinetic assessment in the monkey—The circulation levels of B1344 were examined following the first dose on day 1 and following last dose on day 78 of the whole study. The plasma samples were collected at 0, 4, 8, 12, 24, 48, 72, 120, 168 hours post the first dose and last dose on day 78. The exposure levels of B1344 were determined with the enzyme-linked immunosorbent assays (ELISA) that identifies intact FGF21 specifically.

RNA extraction and real-time quantitative PCR—Mice livers were homogenized and lysed in TRIzol Reagent (Life Technologies) for total RNA extraction according to the manufacturer's protocol. Reverse transcription was then performed with SuperScript II reverse transcriptase (Life Technologies) and Oligo d (T). The resulting cDNA was subjected to real-time PCR with gene-specific primers in the presence of SYBR Green PCR master mix (Applied Biosystems) using StepOnePlus Real-Time PCR System (Applied Biosystems) as described previously^{4, 5}.

Hepatic lipid extraction and measurement—The triglyceride and total cholesterol extraction and measurement from mice livers were described previously. In brief, lipids were extracted from liver homogenates in PBS with chloroform/methanol (2:1), dried under nitrogen gas and then

resuspended in double distilled water. Infinity Triglycerides kit (cat. TR22421) and Total Cholesterol kit (cat. TR13421) obtained from Stanbio Laboratory (Boerne, TX) were used for the measurement of liver lipids. The lipids were normalized to initial liver weight for those of homogenates.

Reference

1. Ye X, Qi J, Sun G, Ren G, Zhu S, Wu Y, Yu D, Zhao J, Liu M, Li D. Enhancement of the pharmacological efficacy of fgf-21 by genetic modification and pegylation. *Curr Pharm Biotechnol.* 2013;14:1287-1298
2. Chalasani N, Younossi Z, Lavine JE, Charlton M, Cusi K, Rinella M, Harrison SA, Brunt EM, Sanyal AJ. The diagnosis and management of nonalcoholic fatty liver disease: Practice guidance from the american association for the study of liver diseases. *Hepatology.* 2018;67:328-357
3. Kleiner DE, Brunt EM, Van Natta M, Behling C, Contos MJ, Cummings OW, Ferrell LD, Liu YC, Torbenson MS, Unalp-Arida A, Yeh M, McCullough AJ, Sanyal AJ, Nonalcoholic Steatohepatitis Clinical Research N. Design and validation of a histological scoring system for nonalcoholic fatty liver disease. *Hepatology.* 2005;41:1313-1321
4. Gong Q, Hu Z, Zhang F, Cui A, Chen X, Jiang H, Gao J, Chen X, Han Y, Liang Q, Ye D, Shi L, Chin YE, Wang Y, Xiao H, Guo F, Liu Y, Zang M, Xu A, Li Y. Fibroblast growth factor 21 improves hepatic insulin sensitivity by inhibiting mammalian target of rapamycin complex 1 in mice. *Hepatology.* 2016;64:425-438
5. Zhang F, Hu Z, Li G, Huo S, Ma F, Cui A, Xue Y, Han Y, Gong Q, Gao J, Bian H, Meng Z, Wu H, Long G, Tan Y, Zhang Y, Lin X, Gao X, Xu A, Li Y. Hepatic crebzf couples insulin to lipogenesis by inhibiting insig activity and contributes to hepatic steatosis in diet-induced insulin-resistant mice. *Hepatology.* 2018;68:1361-1375

Table S2. Summary of Changes in Metabolic parameters and cytokines

	Vehicle	0.5mg/kg BIW	1mg/kg BIW	2mg/kg BIW	2mg/kg QW
Glucose(mg/dl)					
Baseline	177.8±51.3	184.6±89.3	168.3±43.7	145.1±40.6	176.6±45.0
Week 10	196.5±111.3	134.1.0±44.2	132.8±38.2	115.1±49.8	130.4±26.6
Insulin(μUm/l)					
Baseline	261.2±183.0	316.8±307.5	266.3±214.0	231.8±126.1	278.0±218.9
Week 10	286.0±298.0	251.5±396.9	89.9±36.6	81.8±76.4	126.3±74.7
HbA1c(%)					
Baseline	7.0±1.9	6.8±2.4	6.6±1.8	6.4±1.9	6.9±1.8
Week 10	7.7±4.1	5.1±0.6	4.7±1.2	4.7±1.6	4.7±1.3
HbA1c(mmol/mol)					
Baseline	53.0±20.8	51.0±26.2	49.0±19.7	46.0±20.8	52.0±19.7
Week 10	61.0±44.8	32.0±6.6	28.0±13.1	28.0±17.5	28.0±14.2
Fructosamine(μmol/l)					
Baseline	256.0±57.1	285.4±108.5	260.1±56.5	265.7±55.3	278.6±57.0
Week 10	281.3±139.6	241.6±48.9	209.4±27.2	234.1±71.8	231.3±33.7
Triglyceride(mmol/l)					
Baseline	4.6±5.0	5.8±4.8	3.1±2.6	3.3±1.9	3.1±1.0
Week 10	7.7±7.6	2.8±1.7	1.8±2.1	2.9±4.3	1.7±0.8
LDL-C(mmol/l)					
Baseline	10.2±9.8	10.1±6.9	5.0±4.7	10.7±7.3	9.9±7.3
Week 10	9.3±9.1	8.5±6.4	4.8±3.8	11.1±6.0	10.8±4.6
VLDL-C(mmol/l)					
Baseline	2.1±2.3	2.6±2.2	1.4±1.2	1.5±0.9	1.4±0.5
Week 10	3.5±3.5	1.3±0.8	0.8±1.0	1.3±1.9	0.8±0.4
HDL-C(mmol/l)					
Baseline	1.3±0.6	1.3±0.47	1.7±0.9	1.3±0.5	1.3±0.3
Week 10	1.2±0.8	1.5±0.35	1.7±0.68	1.4±0.42	1.5±0.26
Adiponectin(μg/ml)					
Baseline	9.7±6.7	13.8±8.3	18.0±15.9	21.8±7.1	11.1±4.5
Week 10	6.7±3.6	34.8±44.3	25.7±21.8	37.0±22.9	17.5±8.5
Glucagon(pg/ml)					
Baseline	428.3±292.6	496.5±567.5	651.5±611.9	418.7±208.2	472.1±294.2
Week 10	202.2±135.9	424.2±767.7	101.6±75.7	195.0±126.7	459.6±360.8
BUN(mmol/l)					
Baseline	4.0±0.4	5.7±4.8	4.3±0.8	4.1±0.6	3.8±1.1
Week 10	4.2±2.0	4.1±1.4	4.2±0.9	4.5±1.8	3.8±1.0

The table summarizes mean $\pm$ SD values for the metabolic and security parameters in cynomolgus monkeys at baseline and the end of the treatment (week10), n=6-8. Abbreviations: HbA1c, glycated hemoglobin; LDL-C, low-density lipoprotein cholesterol; VLDL-C, very low density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol; BUN, blood urea nitrogen.

Table S3. Statistics from obese cynomolgus monkeys study

Group	Day	Body weight		Triglycerides		VLDL-C	
		mean	P values	mean	P values	mean	P values
Vehicle	8	-0.19		7.21		7.16	
Vehicle	15	-0.75		8.68		8.66	
Vehicle	22	-0.86		27.07		26.92	
Vehicle	29	-1.17		42.55		42.68	
Vehicle	36	-1.60		49.77		49.76	
Vehicle	43	-1.49		50.75		50.81	
Vehicle	50	-1.91		73.96		73.92	
Vehicle	57	-2.05		50.92		50.97	
Vehicle	64	-2.53		42.07		42.72	
Vehicle	71	-2.76					
Vehicle	78	-2.66		69.46		69.57	
Vehicle	85	-1.89					
Vehicle	92	-2.31		24.09		24.27	
Vehicle	99	-2.61					
Vehicle	106	-2.22		34.43		34.50	
B1344 0.5mg/kg biw	8	-1.14	0.0460	-51.24	0.0006	-50.16	0.0006
B1344 0.5mg/kg biw	15	-2.36	0.0336	-48.91	0.0005	-47.52	0.0006
B1344 0.5mg/kg biw	22	-3.49	0.0162	-41.95	0.0008	-40.70	0.0008
B1344 0.5mg/kg biw	29	-4.58	0.0137	-42.00	0.0015	-40.87	0.0015
B1344 0.5mg/kg biw	36	-5.72	0.0194	-28.24	0.0089	-27.56	0.0087
B1344 0.5mg/kg biw	43	-5.65	0.0316	-38.43	0.0040	-37.70	0.0040
B1344 0.5mg/kg biw	50	-7.22	0.8538	-27.01	0.0522	-26.55	0.0528
B1344 0.5mg/kg biw	57	-8.42	0.9366	-27.78	0.0528	-27.07	0.0533
B1344 0.5mg/kg biw	64	-8.83	0.9363	-21.85	0.0570	-21.27	0.0586
B1344 0.5mg/kg biw	71	-9.10	0.0578				
B1344 0.5mg/kg biw	78	-10.16	0.9777	-11.10	0.0242	-10.63	0.0237
B1344 0.5mg/kg biw	85	-10.44	0.8725				
B1344 0.5mg/kg biw	92	-10.23	0.0489	-3.82	0.0931	-2.58	0.0950
B1344 0.5mg/kg biw	99	-9.13	0.0841				
B1344 0.5mg/kg biw	106	-9.48	0.0591	-7.57	0.1009	-16.74	0.0766
B1344 1mg/kg biw	8	-1.01	0.0107	-57.12	<0.0001	-57.12	<0.0001
B1344 1mg/kg biw	15	-2.06	0.0084	-53.77	<0.0001	-53.85	<0.0001
B1344 1mg/kg biw	22	-3.47	0.0008	-46.66	0.0001	-48.71	0.0001
B1344 1mg/kg biw	29	-4.51	0.0005	-52.08	0.0003	-52.24	0.0003
B1344 1mg/kg biw	36	-5.47	0.0004	-43.96	0.0007	-43.94	0.0007
B1344 1mg/kg biw	43	-6.04	0.0006	-55.96	0.0004	-55.97	0.0004
B1344 1mg/kg biw	50	-7.23	0.9552	-51.45	0.0547	-51.57	0.0547
B1344 1mg/kg biw	57	-7.89	0.9185	-43.09	0.0731	-42.89	0.0729
B1344 1mg/kg biw	64	-8.46	0.9110	-29.23	0.0879	-29.19	0.0883

B1344 1mg/kg biw	71	-9.23	0.00018				
B1344 1mg/kg biw	78	-9.63	0.8390	-28.98	0.0016	-28.81	0.0017
B1344 1mg/kg biw	85	-9.47	0.7787				
B1344 1mg/kg biw	92	-9.39	0.0004	14.79	0.3440	14.83	0.3422
B1344 1mg/kg biw	99	-8.05	0.0021				
B1344 1mg/kg biw	106	-8.26	0.0002	38.17	0.2733	77.29	0.3305
B1344 2mg/kg biw	8	-1.40	0.0089	-52.13	0.0014	-52.00	0.0014
B1344 2mg/kg biw	15	-3.38	0.0014	-52.18	0.0003	-52.15	0.0003
B1344 2mg/kg biw	22	-4.39	0.0028	-45.77	0.0006	-46.00	0.0006
B1344 2mg/kg biw	29	-5.74	0.0041	-48.35	0.0008	-48.10	0.0008
B1344 2mg/kg biw	36	-7.13	0.0069	-39.31	0.0024	-39.36	0.0024
B1344 2mg/kg biw	43	-7.53	0.0122	-37.42	0.0051	-37.28	0.0051
B1344 2mg/kg biw	50	-9.00	0.7760	-32.15	0.1781	-31.95	0.2442
B1344 2mg/kg biw	57	-9.95	0.7080	-35.00	0.2407	-34.94	0.3182
B1344 2mg/kg biw	64	-10.83	0.6843	-21.55	0.4384	-21.65	0.5332
B1344 2mg/kg biw	71	-11.89	0.0215				
B1344 2mg/kg biw	78	-12.64	0.5735	-27.75	0.0036	-27.52	0.0034
B1344 2mg/kg biw	85	-12.30	0.5504				
B1344 2mg/kg biw	92	-12.39	0.0238	-5.76	0.5724	-6.06	0.7168
B1344 2mg/kg biw	99	-11.45	0.0370				
B1344 2mg/kg biw	106	-11.55	0.0245	-0.67	0.5282	-0.40	0.6519
B1344 2mg/kg qw	8	-1.20	0.0459	-46.55	0.0013	-46.44	0.0012
B1344 2mg/kg qw	15	-2.00	0.0370	-41.51	0.0015	-41.51	0.0015
B1344 2mg/kg qw	22	-2.71	0.0331	-32.98	0.0013	-33.16	0.0014
B1344 2mg/kg qw	29	-3.85	0.0136	-33.03	0.0035	-33.24	0.0034
B1344 2mg/kg qw	36	-4.76	0.0172	-28.87	0.0058	-28.84	0.0058
B1344 2mg/kg qw	43	-5.33	0.0151	-30.04	0.0053	-30.28	0.0051
B1344 2mg/kg qw	50	-6.63	0.4202	-33.25	0.0512	-33.20	0.0514
B1344 2mg/kg qw	57	-7.23	0.3465	-18.88	0.0693	-18.91	0.0691
B1344 2mg/kg qw	64	-7.84	0.3553	-21.99	0.0720	-21.88	0.0732
B1344 2mg/kg qw	71	-8.33	0.0187				
B1344 2mg/kg qw	78	-8.69	0.2441	-18.13	0.0018	-18.15	0.0017
B1344 2mg/kg qw	85	-9.04	0.1684				
B1344 2mg/kg qw	92	-8.64	0.0065	17.39	0.1480	17.30	0.1486
B1344 2mg/kg qw	99	-7.03	0.0382				
B1344 2mg/kg qw	106	-7.11	0.0105	15.97	0.1178	15.72	0.1176

Group	Day	Adiponectin		LDL-C		HDL-C	
		mean	P values	mean	P values	mean	P values
Vehicle	8			-4.09		0.44	
Vehicle	15			9.71		0.34	
Vehicle	22	4.19		9.06		0.17	
Vehicle	29			-0.46		-1.33	
Vehicle	36			14.77		10.38	
Vehicle	43	-1.01		8.31		6.68	
Vehicle	50			7.64		3.08	
Vehicle	57			-1.98		-2.30	
Vehicle	64			11.69		6.54	
Vehicle	71						
Vehicle	78	-8.97		-26.46		-6.31	
Vehicle	85						
Vehicle	92			-35.26		-3.12	
Vehicle	99						
Vehicle	106	-18.65		-26.17		-2.05	
B1344 0.5mg/kg biw	8			7.45	0.2228	32.36	0.0733
B1344 0.5mg/kg biw	15			15.86	0.7303	32.84	0.1096
B1344 0.5mg/kg biw	22	69.30	0.0403	26.75	0.5054	33.71	0.1049
B1344 0.5mg/kg biw	29			7.93	0.5860	27.77	0.0673
B1344 0.5mg/kg biw	36			6.20	0.3374	35.55	0.2739
B1344 0.5mg/kg biw	43	66.33	0.0389	1.63	0.4441	42.04	0.2358
B1344 0.5mg/kg biw	50			-13.30	0.4610	44.82	0.7887
B1344 0.5mg/kg biw	57			-5.30	0.6134	35.11	0.9589
B1344 0.5mg/kg biw	64			-5.98	0.5352	40.66	0.9618
B1344 0.5mg/kg biw	71						
B1344 0.5mg/kg biw	78	125.58	0.0998	-17.75	0.4835	33.09	0.2760
B1344 0.5mg/kg biw	85						
B1344 0.5mg/kg biw	92			-29.41	0.7184	14.93	0.9668
B1344 0.5mg/kg biw	99						
B1344 0.5mg/kg biw	106	14.32	0.2006	-18.88	0.4478	7.46	0.8780
B1344 1mg/kg biw	8			25.64	0.1318	18.71	0.1754
B1344 1mg/kg biw	15			47.35	0.3043	2.94	0.8436
B1344 1mg/kg biw	22	63.52	0.0043	59.32	0.2224	12.41	0.4880
B1344 1mg/kg biw	29			54.00	0.1778	11.96	0.4597
B1344 1mg/kg biw	36			62.08	0.2065	17.17	0.7041
B1344 1mg/kg biw	43	55.80	0.0063	48.20	0.2807	16.65	0.5805
B1344 1mg/kg biw	50			27.04	0.5387	13.90	0.6327
B1344 1mg/kg biw	57			36.87	0.5542	23.29	0.7526
B1344 1mg/kg biw	64			47.33	0.6225	23.57	0.6614
B1344 1mg/kg biw	71						

B1344 1mg/kg biw	78	47.84	0.0066	14.66	0.1520	11.43	0.4493
B1344 1mg/kg biw	85						
B1344 1mg/kg biw	92			-18.69	0.5362	-1.90	0.4644
B1344 1mg/kg biw	99						
B1344 1mg/kg biw	106	1.12	0.0660	21.73	0.4954	10.29	0.3673
B1344 2mg/kg biw	8			15.36	0.1298	25.58	0.1425
B1344 2mg/kg biw	15			19.11	0.5293	7.70	0.5689
B1344 2mg/kg biw	22	66.79	0.0044	26.16	0.4269	3.34	0.7676
B1344 2mg/kg biw	29			9.78	0.5708	4.72	0.6394
B1344 2mg/kg biw	36			16.70	0.9061	7.08	0.7853
B1344 2mg/kg biw	43	67.87	0.0528	20.11	0.4936	-1.02	0.5638
B1344 2mg/kg biw	50			12.55	0.1696	-2.86	0.4473
B1344 2mg/kg biw	57			16.78	0.2066	3.77	0.5127
B1344 2mg/kg biw	64			19.08	0.1659	7.45	0.4605
B1344 2mg/kg biw	71						
B1344 2mg/kg biw	78	56.23	0.0221	4.82	0.1289	0.05	0.6726
B1344 2mg/kg biw	85						
B1344 2mg/kg biw	92			-29.61	0.2037	-3.78	0.4300
B1344 2mg/kg biw	99						
B1344 2mg/kg biw	106	4.56	0.1830	-19.14	0.2289	4.02	0.4332
B1344 2mg/kg qw	8			22.74	0.0629	18.39	0.1275
B1344 2mg/kg qw	15			19.25	0.6043	13.86	0.2711
B1344 2mg/kg qw	22	68.79	0.0007	28.36	0.3165	19.68	0.1753
B1344 2mg/kg qw	29			31.38	0.0564	24.41	0.1143
B1344 2mg/kg qw	36			40.41	0.1469	28.96	0.1917
B1344 2mg/kg qw	43	67.87	0.0035	32.87	0.1888	21.72	0.3270
B1344 2mg/kg qw	50			7.58	0.1856	22.15	0.6453
B1344 2mg/kg qw	57			15.20	0.2489	21.10	0.5934
B1344 2mg/kg qw	64			20.44	0.2125	23.23	0.5017
B1344 2mg/kg qw	71						
B1344 2mg/kg qw	78	56.23	0.0043	16.61	0.0529	16.62	0.1657
B1344 2mg/kg qw	85						
B1344 2mg/kg qw	92			-20.89	0.2342	8.42	0.4666
B1344 2mg/kg qw	99						
B1344 2mg/kg qw	106	-4.56	0.2802	-9.78	0.2294	10.79	0.4167

Group	Day	FBG		HbA1c		Insulin	
		mean	P values	mean	P values	mean	P values
Vehicle	8	0.45		-4.17		-7.05	
Vehicle	15	-0.12		1.33		-21.41	
Vehicle	22	0.86		1.16		0.04	
Vehicle	29	11.23		-0.18		-24.89	
Vehicle	36	7.61		0.80		-23.81	
Vehicle	43	8.72		-0.97		-19.39	
Vehicle	50	11.84		-4.78		8.96	
Vehicle	57	25.03		3.20		-2.21	
Vehicle	64	17.71		-3.46		-6.31	
Vehicle	71						
Vehicle	78	9.17		2.71		2.54	
Vehicle	85						
Vehicle	92	-3.47		-6.92		-16.84	
Vehicle	99						
Vehicle	106	1.39		0.62		-3.31	
B1344 0.5mg/kg biw	8	-13.40	0.0443	-3.40	0.7212	-24.10	0.3716
B1344 0.5mg/kg biw	15	-12.97	0.0517	-4.82	0.0726	-29.55	0.6122
B1344 0.5mg/kg biw	22	-17.58	0.0563	-8.99	0.0052	-6.84	0.7531
B1344 0.5mg/kg biw	29	-16.14	0.0177	-13.25	0.0099	-14.11	0.4490
B1344 0.5mg/kg biw	36	-13.33	0.0462	-15.83	0.0013	-5.62	0.2527
B1344 0.5mg/kg biw	43	-21.84	0.0052	-21.66	0.0006	-13.14	0.7037
B1344 0.5mg/kg biw	50	-16.67	0.2438	-25.76	0.2218	-11.04	0.4788
B1344 0.5mg/kg biw	57	-22.92	0.0304	-21.68	0.1398	-31.85	0.6236
B1344 0.5mg/kg biw	64	-21.31	0.0564	-27.40	0.1277	-12.38	0.5425
B1344 0.5mg/kg biw	71						
B1344 0.5mg/kg biw	78	-11.32	0.1615	-26.57	0.0026	-11.12	0.5857
B1344 0.5mg/kg biw	85						
B1344 0.5mg/kg biw	92	-16.33	0.6330	-30.27	0.1118	-17.21	0.3615
B1344 0.5mg/kg biw	99						
B1344 0.5mg/kg biw	106	-0.93	0.9326	-22.22	0.1718	14.91	0.4070
B1344 1mg/kg biw	8	-15.99	0.0715	-4.58	0.8817	-39.00	0.0670
B1344 1mg/kg biw	15	-16.59	0.0826	-5.17	0.0772	-41.43	0.1891
B1344 1mg/kg biw	22	-28.03	0.0062	-10.47	0.0089	-36.78	0.0271
B1344 1mg/kg biw	29	-23.92	0.0059	-17.93	0.0018	-57.93	0.0054
B1344 1mg/kg biw	36	-22.01	0.0034	-15.39	0.0024	-45.85	0.0640
B1344 1mg/kg biw	43	-29.84	0.0006	-23.55	0.0012	-41.18	0.1298
B1344 1mg/kg biw	50	-23.89	0.1040	-29.24	0.2323	-50.23	0.5178
B1344 1mg/kg biw	57	-21.32	0.0685	-23.44	0.1805	-29.92	0.8076
B1344 1mg/kg biw	64	-20.53	0.0647	-26.01	0.2611	-29.03	0.8452
B1344 1mg/kg biw	71						

B1344 1mg/kg biw	78	-19.12	0.0737	-27.35	0.0002	-36.18	0.1677
B1344 1mg/kg biw	85						
B1344 1mg/kg biw	92	-3.22	0.9456	-27.17	0.3077	-15.53	0.4497
B1344 1mg/kg biw	99						
B1344 1mg/kg biw	106	0.59	0.9697	-19.56	0.4337	-5.17	0.6746
B1344 2mg/kg biw	8	-17.71	0.0662	-0.76	0.2379	-67.71	0.0044
B1344 2mg/kg biw	15	-15.60	0.1145	-3.31	0.1990	-63.26	0.0074
B1344 2mg/kg biw	22	-21.88	0.0480	-9.17	0.0166	-51.74	0.0049
B1344 2mg/kg biw	29	-16.05	0.0257	-15.61	0.0048	-54.48	0.0619
B1344 2mg/kg biw	36	-11.79	0.1076	-13.43	0.0052	-50.60	0.0917
B1344 2mg/kg biw	43	-23.55	0.0092	-16.78	0.0297	-47.26	0.1669
B1344 2mg/kg biw	50	-14.76	0.2212	-22.13	0.5662	-49.23	0.4486
B1344 2mg/kg biw	57	-15.83	0.0997	-19.85	0.3906	-54.59	0.2823
B1344 2mg/kg biw	64	-15.16	0.1715	-22.29	0.5452	-51.62	0.3274
B1344 2mg/kg biw	71						
B1344 2mg/kg biw	78	-19.47	0.0650	-24.28	0.0072	-55.94	0.0326
B1344 2mg/kg biw	85						
B1344 2mg/kg biw	92	-14.42	0.5687	-21.24	0.7282	-34.94	0.9950
B1344 2mg/kg biw	99						
B1344 2mg/kg biw	106	-7.86	0.5400	-21.50	0.4616	-20.66	0.8344
B1344 2mg/kg qw	8	-3.65	0.5413	-0.89	0.2964	-39.59	0.0920
B1344 2mg/kg qw	15	-10.83	0.0742	-1.21	0.4511	-27.63	0.6253
B1344 2mg/kg qw	22	-12.49	0.0921	-8.12	0.0275	-31.75	0.0223
B1344 2mg/kg qw	29	-11.70	0.0051	-13.47	0.0189	-42.33	0.0752
B1344 2mg/kg qw	36	-11.92	0.0386	-13.41	0.0018	-43.43	0.0887
B1344 2mg/kg qw	43	-13.71	0.0316	-20.20	0.0044	-27.31	0.6245
B1344 2mg/kg qw	50	-10.29	0.1485	-23.56	0.1770	-24.50	0.3819
B1344 2mg/kg qw	57	-12.26	0.0653	-25.01	0.0625	-34.94	0.2196
B1344 2mg/kg qw	64	-17.05	0.0623	-26.52	0.0992	-30.72	0.2974
B1344 2mg/kg qw	71						
B1344 2mg/kg qw	78	-14.72	0.0980	-27.27	0.0025	-38.98	0.0747
B1344 2mg/kg qw	85						
B1344 2mg/kg qw	92	-13.16	0.4323	-21.00	0.3530	-11.03	0.9867
B1344 2mg/kg qw	99						
B1344 2mg/kg qw	106	-2.18	0.5781	-20.82	0.1435	12.48	0.7312

Abbreviations: VLDL-C, very low density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol; FBG, fasting blood glucose; HbA1c, glycated hemoglobin.