

Checklist for Reporting Human Islet Preparations Used in Research

Adapted from Hart NJ, Powers AC (2018) Progress, challenges, and suggestions for using human islets to understand islet biology and human diabetes. Diabetologia <u>https://doi.org/10.1007/s00125-018-4772-2</u>.

Manuscript DOI: https://doi.org/10.2337/[insert manuscript submission number] (Example, https://doi.org/10.2337/db18-1234)

Title: Potential protection against type 2 diabetes in obesity through lower CD36 expression and improved exocytosis in β -cells

Author list: Mototsugu Nagao, Jonathan L.S. Esguerra, Akira Asai, Jones K. Ofori, Anna Edlund, Anna Wendt, Hitoshi Sugihara, Claes B. Wollheim, Shinichi Oikawa, Lena Eliasson

Corresponding author: Lena Eliasson, Mototsugu Nagao

Email address: lena.eliasson@med.lu.se, s8067@nms.ac.jp

Islet preparation	64	66	67	68	69	70	71	72ª
MANDATORY INFORMATION								
Unique identifier	144	147	150	153	154	160	161	162
Donor age (years)	32	66	69	47	64	74	57	57
Donor sex (M/F)	М	М	F	F	М	F	F	М
Donor BMI (kg/m ²)	24.7	23.1	19.5	34.6	24.7	23.9	38.1	23.4

Donor HbA1c (%)	5.4	5.6	5.8	7	5.7	6.3	5.5	6
Origin/source of islets ^b	NICS							
Islet isolation centre	Uppsala University							
Donor history of diabetes? Yes/No	No	No	No	Yes	No	Yes	No	No
If Yes, complete the next two lines if this information is available								
Diabetes duration (years)								
Glucose-lowering therapy at time of death ^c								

RECOMMENDED INFORMATION								
Donor cause of death								
Warm ischaemia time (h)								
Cold ischaemia time (h)								

Estimated purity (%)	37	65	68	43	89	82	85	53
Estimated viability (%)								
Total culture time (h) ^d								
Glucose-stimulated insulin secretion (Stimulation index) ^e	5.1	3.6	8.8	3.1	1.7	21.2	12	8.7
Handpicked to purity? Yes/No								
Additional notes								

^aIf you have used more than eight islet preparations, please complete additional forms as necessary
^bFor example, IIDP, ECIT, Alberta IsletCore
^cPlease specify the therapy/therapies
^dTime of islet culture at the isolation centre, during shipment and at the receiving laboratory
^ePlease specify the test and the results: Stimulation index was calculated by dividing the average insulin concentration of the high-glucose phase by that of the low-glucose phase in a dynamic perfusion.