

Supplemental table 1. Hazard ratios for all-cause and cardiovascular mortality according to different levels of HbA1c

	HbA1c (%)						
	<6	6-6.9	7.0-7.9	8.0-8.9	9.0-9.9	10-10.9	≥11
All-cause mortality							
No. of participants	293	990	1156	998	878	709	1087
No. of death	33	83	128	156	103	111	201
Person-years	2002.9	7127.6	7992.8	6873.5	6168.1	5022.6	8035.3
Adjusted HR (95% CI)							
Model 1	1.36 (0.92-2.02)	1.00	1.22 (0.95-1.58)	1.82 (1.42-2.33)	1.36 (1.04-1.79)	1.87 (1.43-2.44)	2.06 (1.63-2.61)
Model 2	1.53 (1.02-2.31)	1.00	1.23 (0.94-1.60)	1.65 (1.27-2.13)	1.19 (0.89-1.59)	1.65 (1.25-2.19)	1.89 (1.47-2.45)
Cardiovascular mortality							
No. of death	14	21	46	58	43	37	58
Adjusted HR (95% CI)							
Model 1	2.06 (1.10-3.88)	1.00	1.55 (0.98-2.45)	2.41 (1.56-3.73)	1.98 (1.25-3.14)	2.17 (1.34-3.5)	2.07 (1.34-3.21)
Model 2	1.91 (0.97-3.78)	1.00	1.59 (0.99-2.53)	2.32 (1.47-3.67)	2.00 (1.23-3.25)	2.09 (1.26-3.45)	2.25 (1.40-3.61)

HR, hazard ratio; CI, confidence interval.

Model 1 adjusted for age and sex; Model 2 adjusted for age, sex, smoking, diabetes duration, body mass index, systolic blood pressure, triglycerides, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, history of cancer, history of cardiovascular diseases, using antihypertensive drugs, aspirin and statins.