Figure S1. Patient education handout utilized by CPS

Diabetes increases your risk of heart attack and stroke

How can statins lower my risk?

- Diabetes is one of the major risk factors for heart disease
- Statins are prescription medications used to lower cholesterol levels in individuals with heart disease or those at increased risk of developing heart disease or stroke
- Common statins include rosuvastatin, atorvastatin, simvastatin, and pravastatin
- Statins have been shown to lower your chance of heart disease regardless of your cholesterol levels.
- It is important to continue taking your statin medication even if you feel well
- Do not stop taking this medication without talking to your doctor or pharmacist
- Always check with your doctor or pharmacist before starting a new medication, including overthe-counter medicines and vitamins/herbal supplements

Call your doctor right away if you of the following after starting a statin:

- Have unexplained muscle pain or weakness
- Notice your urine (pee) is dark brown or coca-colored
- Notice your skin or the whites of your eyes become yellow
- Have very bad stomach pain or persistent vomiting
- Have a skin rash

How can I refill my prescription?

- 1. Call automated refill phone line
- 2. Online at www.myhealth.va.gov
- 3. Mail refill slip to the pharmacy
- 4. Place refill slip in mail slot in the pharmacy



Image from Microsoft Word 2002

Figure S2. Incorporating the process into the current workflow of clinical pharmacist and potential utilization of clinical pharmacy technician for reviewing chart review components



