

Supplemental Table 1. ApoM-induced Risk Ratios at DCCT Baseline and Closeout stratified by disease status and DCCT Randomized Treatment group.

Measure	Disease Progression (groups are not mutually exclusive)		
	Persistently Normal	Macroalbuminuria	Chronic Kidney Disease
INTENSIVE TREATMENT ARM	(n=180)	(n=12)	(n=7)
Overall	--	1.63 (1.03,2.59)	2.15 (1.10, 4.19)
Baseline DCCT	--	1.28 (0.68,2.39)	1.59 (0.94,2.69)
End of DCCT	--	2.19 (1.14,4.19)	2.75 (1.05,7.22)
STANDARD TREATMENT ARM	(n=149)	(n=38)	(n=19)
Overall	--	1.01 (0.85,1.20)	1.31 (1.00, 1.73)
Baseline DCCT	--	1.03 (0.80,1.31)	1.29 (0.79,2.13)
End of DCCT	--	1.01 (0.80,1.29)	1.35 (0.96,1.90)

Risk Ratios are adjusted for baseline disease cohort, sex, triglycerides, co measured eGFR, AER and the use of any ACE/ARB medications prior to disease progression.

Risk ratios are for a 1 standardized unit increase in ApoM and 95% confidence interval compared to persistently normal. The standard unit is measured in terms of a z-score.